

CUSTOMER TREATMENT & SAFETY NOTICE



Client Details

Client Name: _____
Site Address: _____
Date of Treatment: _____
Technician: _____

Treatment Information. Pest Treated

- Rats
- Mice
- Wasps
- Cockroaches
- Fleas
- Ants
- Other: _____

Areas Treated

Product Information

Product Used: _____
Active Ingredient: _____
PCS/HSE No: _____

Method of Application:

- Baiting
- Residual Spray
- Gel
- Dust
- ULV/Fogging
- Monitoring Device
- Other: _____

Quantity/Concentration Used:

Important Safety Information

- Do not touch treated areas or bait stations
- Keep children and pets away from treatment areas
- Do not wash treated surfaces unless advised
- Ventilate rooms after treatment if instructed
- Report any concerns immediately

Emergency Advice In case of accidental exposure:

- Wash affected skin thoroughly
- Rinse eyes with clean water
- Seek medical advice if symptoms occur
- Have product information available when seeking treatment

Technician Recommendations

Emergency Contact - _____

Technician Signature: _____

Client Signature (optional): _____

CUSTOMER TREATMENT & SAFETY NOTICE



Rodent Treatment (Rodenticide) Advice & Safety Information

Treatment Advice

Rodenticides have been used as a last resort under Integrated Pest Management (IPM) principles at your property. Rodents may take 7–10 days to consume a lethal dose of bait. A follow-up visit will normally be carried out approximately 14 days after the initial treatment to replenish bait and assess activity levels. In many cases, two visits are sufficient; however, larger or established infestations may require additional visits and treatments. For your safety and to comply with current legislation, bait stations, traps, and treatment points must not be interfered with, moved, or tampered with.

Important Safety Warning

Tamper-resistant bait stations have been placed at the property as part of the treatment programme. These bait stations must not be touched, opened, or moved. It is the responsibility of the property owner/occupier to ensure that children and pets are kept away from all treatment areas and bait points at all times.

Rodenticide products used may contain:

- Bromadiolone
- Difenacoum

If accidental poisoning or exposure is suspected, seek immediate medical attention and provide details of the product used.

COSHH Information

This treatment information and relevant safety advice have also been provided by email in accordance with the Control of Substances Hazardous to Health Regulations (Northern Ireland) 2003 (COSHH). Safety Data Sheets (SDS) and COSHH assessments are available upon request.

After Treatment Recommendations

Following successful eradication, proofing measures should be carried out to minimise future infestations:

1. Seal gaps around pipes, vents, doors, and outbuildings.
 - Rats require gaps of approximately 15mm
 - Mice require gaps of approximately 6mm
2. Keep gardens, sheds, and storage areas clean and free from debris, vegetation, and wood piles.
3. Ensure drain covers and inspection chambers are correctly fitted and maintained.
4. Store waste securely with closed bin lids and covered compost areas.
5. Avoid excessive bird feeding and minimise food spillage by using bird feeders or tables.

Thank you for choosing District Pest Control.